

Contents lists available at [SciVerse ScienceDirect](http://www.sciencedirect.com)

Cancer Treatment Reviews

journal homepage: www.elsevierhealth.com/journals/ctrv

Laboratory-Clinic Interface

Taxane resistance in breast cancer: Mechanisms, predictive biomarkers and circumvention strategies

S. Murray^{a,*}, E. Briasoulis^b, H. Linardou^c, D. Bafaloukos^b, C. Papadimitriou^d^a Department of Molecular Oncology, GeneKOR, Athens, Greece^b Cancer Biobank Center, University of Ioannina Medical School, Ioannina, Greece^c 1st Department of Medical Oncology, Metropolitan Hospital, Athens, Greece^d Department of Clinical Therapeutics, University of Athens School of Medicine, Alexandra Hospital, Athens, Greece

ARTICLE INFO

Article history:

Received 17 September 2011

Received in revised form 15 February 2012

Accepted 24 February 2012

Available online xxxx

Keywords:

Stathmin
P-glycoprotein
Resistance
Taxane
p53
HER2
BRCA1/2
MRP-1

ABSTRACT

Background: Taxanes are established in the treatment of metastatic breast cancer (MBC) and early breast cancer (EBC) as potent chemotherapy agents. However, their therapeutic usefulness is limited by *de-novo* refractoriness or acquired resistance, which are common drawbacks to most anti-cancer cytotoxics. Considering that the taxanes will remain principle chemotherapeutic agents for the treatment of breast cancer, we reviewed known mechanisms of resistance in with an outlook of optimizing their clinical use.

Methods: We searched the PubMed and MEDLINE databases for articles (from inception through to 9th January 2012; last search 10/01/2012) and journals known to publish information relevant to taxane chemotherapy. We imposed no language restrictions. Search terms included: cancer, breast cancer, response, resistance, taxane, paclitaxel, docetaxel, taxol. Due to the possibility of alternative mechanisms of resistance all combination chemotherapy treated data sets were removed from our overview.

Results: Over-expression of the MDR-1 gene product Pgp was extensively studied *in vitro* in association with taxane resistance, but data are conflicting. Similarly, the target components microtubules, which are thought to mediate refractoriness through alterations of the expression pattern of tubulins or microtubule associated proteins and the expression of alternative tubulin isoforms, failed to confirm such associations. Little consensus has been generated for reported associations between taxane-sensitivity and mutated p53, or taxane-resistance and overexpression of Bcl-2, Bcl-xL or NFkB. In contrary sufficient *in vitro* data support an association of spindle assembly checkpoint (SAC) defects with resistance. Clinical data have been limited and inconsistent, which relate to the variety of methods used, lack of standardization of cut-offs for quantitation, differences in clinical endpoints measured and in methods of tissue collection preparation and storage, and study/patient heterogeneity. The most prominent finding is that pharmaceutical down-regulation of HER-2 appears to reverse the taxane resistance.

Conclusions: Currently no valid practical biomarkers exist that can predict resistance to the taxanes in breast cancer supporting the principle of individualized cancer therapy. The incorporation of several biomarker analyses into prospectively designed studies in this setting are needed.

© 2012 Elsevier Ltd. All rights reserved.

Introduction

Breast cancer remains the most common type of cancer in women, with more than one million reported new cases diagnosed per year.¹ Of those, 20–30% present with metastatic or locally advanced disease, and other 30% will develop recurrent or metastatic disease.² Treatment options include surgery, radiotherapy and systemic treatment. Among the most commonly used cytotoxic drugs for breast cancer are the taxanes; paclitaxel and docetaxel.³

Taxanes were first introduced into clinical use during the 1990's. Both, paclitaxel and docetaxel compared favorable by terms of efficacy in metastatic breast cancer (MBC) and early stage breast cancer (EBC) when tested against older drugs.^{4–7} Today, both taxanes have been established as a viable option in the treatment of MBC and have been incorporated into the management of EBC in association with anthracyclins and trastuzumab where and when appropriate.^{3,8–10}

Although improvements have been made, for virtually all therapeutic strategies, many patients have and eventually almost all patients will develop tumors that are non-responsive to our current treatment strategies whether they are of the so called 'targeted' or 'non-targeted' class.¹¹ Efforts to move more and more patients into adjuvant based therapeutic strategies has highlighted

* Corresponding author. Address: GeneKor A.E., 52 Spaton Ave, Gerakas, 15433 Athens, Greece. Tel.: +30 210 603 2138, mobile: +30 6944644281; fax: +30 210 603 2148.

E-mail address: smgenedb@gmail.com (S. Murray).

the need to either identify those that are less likely to be resistant and/or to develop strategies to circumvent such resistance mechanisms.^{12,13}

In a more simplified view resistance can be *de-novo* (inherent insensitivity) or acquired (due to the emergence of resistant populations). The development of tumor resistance (acquired) is potentially a result of several alterations in the tumor including but not limited to protein isoform switching/dysregulation/mutations; alterations in drug efflux mechanisms, apoptotic modulation, and a number of other candidate mechanisms have been suggested.^{14,15}

One of the most often studied mechanisms with regard to taxane resistance has centered on that of *de-novo* and acquired resistance with respect to drug efflux proteins. These are an ever-enlarging family of proteins that are known to limit drug efficacy by removal at their site of action. These proteins clear excessive extra- and/or intra-cellular concentrations of a variety of substrates and toxins. It is now well known that various cancer cell types express proteins of the adenosine triphosphate (ATP)-binding cassette (ABC) transporter family. The most well known member of the family is the P-glycoprotein (P-gp) membrane protein encoded by the *MDR1* gene, and other similarly functional transporters that have been correlated with reduced efficacy of a variety of different chemotherapeutics, including the taxanes.^{16,17}

Circumvention or blocking resistance mediated by these mechanisms has been both a therapeutic target, but also a clinical challenge.¹⁸ The synthesis of low susceptibility to resistance mechanism analogs of several chemotherapeutic agents has been a continual process.¹⁷ Furthermore, several small molecule inhibitors of Pgp and MRP1 have entered clinical development, unfortunately with limited success.^{19–21} Other strategies have included the development of alternative forms of taxanes that are poor substrates to Pgp, the most clinically advanced of which are the epithiliones.^{22,23}

Whatever agents or strategies we develop will ultimately depend upon our understanding of the mechanism of action of each of the therapeutic agents we develop and administer to our patients. Classification of all patients tumors based upon several measures will hopefully achieve this goal.^{24–30} Considering that the taxanes will remain a principle chemotherapeutic agent for the treatment of breast cancer, a rational understanding not only of predictors of response but also potential predictors of resistance (*de-novo* or acquired) may assist in this personalized approach. This review aims to document our current best evidence regarding mechanisms of taxane resistance and propose potential avenues for circumvention.

Research methodology

The information for this review was obtained by searching the PubMed and MEDLINE databases for articles published until 9th January 2012 (last search 10/01/2011). Electronic early-release publications were also included. We searched journals known to publish information relevant to our topic and cross-referenced the reference lists of recovered articles. We did not impose language restrictions. Search terms included: cancer, breast cancer, response, resistance, taxane, paclitaxel, docetaxel, taxol. Cell line and other *in-vitro* data have been used for mechanistic descriptions; however, precedence has been given to clinical evidence. Data was limited by treatment or pre-treatment strategies, where-in data included in the tables are derived solely from studies with taxane resistant populations or single agent taxane treated populations, i.e. studies with polychemotherapy inclusive of a taxane have not been included due to unknown characterization of ‘other’ agent(s) effect on resistance. Due to the possibility of alternative

mechanisms of resistance all combination chemotherapy treated data sets were removed from our overview. Observations conveyed to the authors by personal communication and unpublished observations were also included. We also contacted experts in the field to broaden our yield of potentially eligible articles. Studies published exclusively in abstract form were not considered (they were considered open to subsequent modification).

The taxanes

Paclitaxel is a plant derivative of the Pacific Yew (*Taxus brevifolia*) and a potent cytotoxic microtubule-stabilizing agent.³¹ It has been found to be efficacious in the treatment of a number of human cancers including ovarian cancer, breast cancer, NSCLC, and other malignancies.^{32–38} However, it has become obvious that many patients treated with paclitaxel present *de-novo* or will acquire resistance to this agent.

Docetaxel is regarded as a second-generation taxane. It is semi-synthetically derived from the esterification of a side chain to 10-deacetyl-beccatin III.³⁹ The chemical status of the two taxanes is almost identical. Docetaxel is typically administered in a vehicle with low hypersensitivity. The both share similar, but not identical, pharmacokinetics and related side effects.⁴⁰ Reported mechanisms of resistance are typically if not identical for both.

A list of some of the taxane formulations available in clinical practice or under investigation is shown in Table 1.

Mechanisms of taxane action

Classically taxanes exert their action through binding to β -tubulin, components of microtubules resulting in the formation of stable microtubules.^{39,41–43} Subsequent arrest at the mitotic checkpoint results in apoptosis presumably through G₂/M arrest and subsequent apoptosis through the mitochondrial pathway.^{44–46} Paclitaxel can also cause disruption of microtubules during interphase, thereby disrupting growth and metabolism.

Some of the most well characterized mechanisms of molecular action include (Fig. 1):

- (a) activation of cell division control-2 kinase (cdc-2),⁴⁷
- (b) stabilization of cyclin B-1,^{48,49}
- (c) activation of the spindle assembly checkpoint,⁵⁰
- (d) induction of apoptosis through phosphorylation of bcl-2,^{51,52}
- (e) inhibition of cell proliferation.⁵³

Both the dose and the duration of exposure appear to be important in triggering apoptosis. At sub-nanomolar concentrations paclitaxel favors microtubule assembly through reduction of the critical concentration of tubulin dimmers, GTP and microtubule associated proteins (MAPs).^{54,55} At picomolar concentrations it also

Table 1
Formulations of taxanes for cancer therapy.

Cationic PEGylated liposomal paclitaxel (EndoTAG-1)
Docetaxel (Taxotere®)
Liposomal docetaxel (ATI-1123 PSN™)
Liposomal docetaxel (ThermoDox®)
Liposomal paclitaxel (LipoTaxen™)
Nanoparticle albumin-bound (NAB) paclitaxel (Abraxane®)
OncoGel (a biocompatible, biodegradable, controlled release depot formulation of paclitaxel in ReGel)
Paclitaxel (Taxol®)
Paclitaxel poliglumex (CT-2103), paclitaxel linked to a biodegradable polyglutamate polymer (OPAXIO™)
Vitamin E based paclitaxel emulsion (Tocosol®)

Incomplete list of several of the most widely known taxane formulations.




Fig. 1. Regulation of cell cycle in relation to taxane resistance. Cell cycle and phase commitment. Mitogenic stimuli converge to activate cyclin D complexes; these allow E2F to activate the expression of genes required for S-phase entry in a pRb dependent fashion. Antiproliferative signals, by means of as yet unclear mechanisms, affect p27^{Kip1} that in turn antagonizes cyclin E-CDK2 activity. The cyclin dependent kinases (cdk1 (cdc2), cdk2, cdk4/6) bind to and regulate cyclin function. Phosphorylation of the complexes regulates various transcriptional events in cell cycle progression. The spindle assembly checkpoint (SAC) appears to be the principal site for taxane induced cell death signals to take effect. As indicated in the text, taxanes have been correlated with differential regulation of a number of key genes (proteins) associated with the cell cycle. Many of these are considered to specifically regulate the S/G2 transition preceding M. Resistance to taxanes is thought to occur through a variety of mechanisms, many of which have been linked to defects in the SAC. Several chemotherapeutic agents act in distinct phases of the cell cycle: G1 (alkylating agents, platinum, cytotoxic antibiotics), S (anti-metabolites, topoisomerase inhibitors, docetaxel), and M (paclitaxel, docetaxel, epothilones, vinca alkaloids). CDK, cyclin-dependent kinase; Cip, CDK interacting protein; HDAC, histone deacetylase; Ink4, Inhibitor of CDK4; Kip, Kinase inhibitory protein, PP-I, protein phosphatase 1; pRb, product of the retinoblastoma susceptibility gene. *Cell cycle checkpoint.

exerts its effects on interphase microtubules and genes controlling apoptosis.⁵⁶ Increasing concentrations result in shifting the equilibrium of dimers to polymers thereby preventing disassociation even under conditions of extreme stress.^{41,57} As a result of the increasing number of patients being treated with taxanes the development of taxane resistance is becoming a clinically important issue. Therefore, the elucidations of resistance are not only important for the development of strategies to overcome it, but also in possibly predicting response of patients to taxane based regimens.

Molecular mechanisms of taxane resistance

Although numerous mechanisms of drug resistance have been recognized we focus on those specifically reporting on taxane resistance. Several mechanisms have been identified in breast cancer cell lines, while characterization of resistance has proven more difficult in clinical specimens. We primarily report on *in vivo* data and supplement with *in vitro* data for a number of the best-characterized mechanisms.

P-glycoprotein (Pgp)

A feature common to most cancer types is multi-drug resistance, i.e. cross resistance of cancer cells to structurally unrelated cytotoxic agents.^{58,59} Several mechanisms of variable drug and

cancer specificity have been associated with the study of cancer cells to cytotoxic xenobiotics.^{60–64}

One of the most well known mechanisms relates to drug resistance associated with the over-expression of the MRD-1 gene product Pgp (permeability-glycoprotein) in cancer.^{65–67} Increased expression (as assessed by immunohistochemistry) of this protein has been extensively studied in taxane resistant breast carcinomas.⁶⁸ However, it remains an elusive marker for clinical implementation due to conflicting data and a lack of standardization in light of consensus recommendations that date back to the mid 1990's.⁶⁹ Pgp is a member of a growing family of at least 49 adenosine triphosphate (ATP) binding cassette (ABC) transporters.^{70,71} Proteins in this family include Pgp/ABCB1, breast cancer resistance protein (BCRP)/ABCG2 and multi-drug resistance related protein (MRP-1)/ABCC1, all of which confer an MDR phenotype.^{72–76} The family is broken down into eight subgroups, Table 2, with each having a general structure as indicated in Fig. 2.

The gene for Pgp is the most widely studied of all resistance mechanisms in breast cancer.^{77–79} It is localized to chromosome 7, encoding a 170 kDa protein containing two ATP-binding sites and two transmembrane domains.⁸⁰ Pgp expression is correlated with acquired and *de-novo* resistance to natural amphipathic products including taxanes, vinca alkaloids, epipodophylotoxins and anthracyclines, confounding separation of taxane specific resistance in light of most therapeutic schedules being polychemotherapy based.⁶⁵

Table 2
ABC transporter subfamilies.

Family	Members	Function	Examples
ABCA ABCB	Largest family, 5 located on 17q24 Consists of 4 full and 7 half transporters	Transport cholesterol and lipids. Located in blood–brain barrier, liver, mitochondria. Transport peptides and bile.	ABCA1 (ABC-1), ABCA12 ABCB5, ABCB1 (MDR-1)
ABCC	Consists of 12 full transporters	Function in ion transport, as cell surface receptor and excrete toxins.	ABCC1 (MRP1), ABCC2 (MRP2), ABCC3 (MRP3)
ABCD ABCE/ABCF	Consists of 4 half transporters Consists of ABCE and 3 × ABCF transporters	All located in peroxisomes These are ATP binding domains derived from the family lacking transmembrane domains. They regulate protein synthesis and expression.	ABCD1 (ALD) ABCE, ABCF1 (ABC27)
ABCG	Consists of 6 reverse half transporters	Transports lipids, drug substrates, bile cholesterol and steroids	ABCG1, ABCG2 (BCRP)

There are 51 known ABC transporters known in humans, classified into 7 families; several synonyms for each exist; however the preferred terminology is derived from the HGCN (HUGO Gene Nomenclature Committee).




Fig. 2. Cartoon structure of ABC transporters. There are currently 51 different human ABC transporters classified into 7 families (Table 2). Due to the vast variety of such transporters as indicated in the Table, we include a cartoon of the ABCG family that is the most representative of data pertaining to taxane resistance. Note here that not all transporters have similar structures. This family currently includes 8 members (1–8). ABCG2 is also known as ABCP and BCRP1 which is known to be expressed in the breast. Typical structure consists of 6 transmembrane beta-barrels; both N- and C- termini are intracellular; a 12 amino acid stretch confers family divergence.

Tubulin

Microtubules, the target components of taxanes, are composed of a backbone of tubulin heterodimers consisting of α - and β -tubulin subunits that combine stoichiometrically to form tubulin dimers in association with microtubule associated proteins (MAPs). The integrity of these tubules is essential for the separation and segregation of chromosomes during cell division wherein microtubule dynamics are critical for the proper alignment of chromosomes and kinetochores, the movement of chromosomes during metaphase and their segregation in anaphase and telophase. They are also involved in development and maintenance of cell shape, the transport of vesicles, mitochondria and other cellular components, and are also involved in signaling.^{81–83}

Throughout the cell cycle levels of tubulin heterodimers and polymerized microtubules are considered to be highly dynamic.⁸⁴ Polymerization is influenced by a number of factors such as GTP (binds to one exchangeable site on β -tubulin and on one non-exchangeable site on α -tubulin), and MAPs. MAPs in themselves constitute a complex family of proteins including MAP2, MAP4, Mip-90, tau and STOP.⁸⁵ Both ends of the microtubules are in a dynamic flux of what are called positive and negative states, wherein the microtubules often appear to treadmill (i.e. elongate, grow, move).⁸⁶ (Fig. 3)

Currently several members of the tubulin superfamily have been identified.⁸⁷ Of them α - and β -tubulin are structurally quite similar (Fig. 4), and each has three functional domains. Tubulin γ -tubulin exists but it is less abundant and is principally associated with the centromeres.^{88,89} The paclitaxel binding site lies in the so called intermediate domain of β -tubulin.⁹⁰ Taxanes, however, only bind to polymerized tubulin, as apposed to other tubulin poisons colchicine and vinca alkaloids that bind to soluble tubulin, therein altering the on and off rate constant of polymerization-depolymerization.⁴¹ In the presence of the taxanes, hydrolysis of GTP to GDP occurs but subsequent depolymerisation is prevented. In the presence of purified tubulin, lateral polymerization and promotion of microtubule stability (and bundle formation) are favored.⁹⁰ In doing so these agents have been termed “microtubule stabilizing agents”, and in their presence cells are conditionally locked into G₂/M, that typically results in death by apoptosis.

Data supporting the potential mechanisms of taxane resistance are reported to include:

- alteration of the expression pattern of α - and β -tubulin in various cell lines⁹¹
- increased expression of tubulin per se⁹²
- the expression of alternative tubulin isoforms⁹³
- alterations in the expression profile of MAPs⁹⁴

There are, however studies that failed to confirm such associations.⁹⁵ Still our understanding of taxane-microtubule interaction(s) remains relatively naïve as we understand that microtubules are involved in numerous cellular functions,⁹⁶ therein the effects of taxanes may be complicated by our snap-shot view of cellular processes leading to cell death.

β -Tubulin isotypes

Theoretically there are differences in paclitaxel binding dependent upon the tubulin isotype, suggesting that these may be important in terms of resistance.⁹⁷ Generally class III appears to be unique in that it destabilizes microtubules and thus much work has concentrated on taxane resistance associated with this isotype. This isoform along with that of isotype I are regarded as the most important for breast cancer as they are potentially clinically useful predictors of response to taxanes (Table 3, Fig. 4). To date, however, the vast majority of data derives from *in vitro* studies, and there are contradictory results between groups and differences between tumor types.^{98,99} Furthermore, most of the data available are derived from experiments in which taxanes are used at concentrations above that clinically available, and/or of much longer durations of exposure, and thus these data may not necessarily reflect the clinical setting.




Fig. 3. Dynamic state of polymerization and depolymerization of microtubules. Microtubules form when heterodimers of a 1:1 ratio of α - and β -tubulin units associate in a head to tail fashion. Cylindrical microtubules consist of 13 proto-filaments with what are termed a plus (+) and minus (–) end. Tubes form and elongate at the plus end when GTP is hydrolyzed to GDP+Pi (inorganic phosphate) that generates a stable GTP cap. At the minus end the off rate is almost equivalent to the on rate, however, there tends to be more rapid elongation at the plus end. The motion of growth at the plus end has been termed ‘treadmilling’ wherein the plus end continues to grow and the minus end slowly recedes. The GTP cap stabilizes the plus end, allowing rapid elongation; this end is also further stabilized by the binding of paclitaxel or docetaxel promoting polymerization. The vinca alkaloids cause destabilization of the GTP/GDP+Pi cap and thus an increase in depolymerization. Colchicine binds at a domain located between the α - and β -subunits suppressing microtubule dynamics. The taxanes bind to the inner surface of the proto-filament while the vinca alkaloids bind to the outer face.




Fig. 4. Alpha and beta tubulin structure and isotypes. There are at least 6 α -tubulin and 8 β -tubulin (I, II, III, IVa, IVb, V, VI, VII) isotypes, and each isotype is different at the amino acid level and in their expression patterns.⁷⁰ Isotypes β I and β IVb are constitutively expressed in all tissues while classes II, III, IVa are typically expressed in neuronal tissues. β VI expression is restricted to cells of blood lineage and hematopoietic tissues.⁷⁶ β III is somewhat different from the others in amino acid sequence and posttranslational modifications. Both β III and β V over-expression have been linked to taxane resistance.^{77,78} The N-terminal domain (Rossmann fold) consists of 6 alternating parallel β -sheets and alpha helices that are involved in nucleotide binding (GDP/GTP). Amino acid residues 207–384 form the lateral and longitudinal constant regions between the α - and β -tubulins in the heterodimer. The c-terminal domain(s) contain two anti-parallel helices (starting at amino acid 385) responsible for isoform specific binding to microtubule associated proteins (inclusive of tau and stathmin).

Mutations of β -tubulin

The earliest report of somatic mutations in β -tubulin was in the amino acid stretch 250–300. Subsequent studies have reported mutations at nucleotides 810 and 1092 of the HM40 isotype of

β -tubulin in paclitaxel resistant cell lines.⁹⁵ In order to explain why mutations occurring outside of the paclitaxel-binding domain correlate with resistance authors have speculated that they lead to alterations in microtubule dynamics. Little data exists regarding breast cancer except for a reported germ line polymorphism at

Table 3*In vivo* data correlating taxane resistance to molecular markers.

Factor	Patient No	Patient Type	Treatment	Response	Comment	References
Bcl-2	63	Neo-adjuvant primary	Docetaxel	Low: 12R vs 34NR High: 3R vs 13NR NS	IHC	[172]
	63	Primary or recurrent	Docetaxel	Bcl-2 ⁺ : 12R vs 5NR Bcl-2 ⁻ : 23R vs 23NR <i>P</i> = 0.144	IHC	[104]
	92	Primary	Docetaxel	Bcl-2 ⁺ : 10R vs 39NR Bcl-2 ⁻ : 7R vs 36NR <i>P</i> = 0.611	IHC Response by pCR (pathological complete response)	[173]
BRCA1	61	Primary or recurrent	Docetaxel	BRCA1 ⁺ : 25R vs 22NR BRCA1 ⁻ : 8R vs 6NR <i>P</i> = 0.794	IHC	[104]
	63	Neo-adjuvant primary	Docetaxel	Low: 5R vs 9NR High: 9R vs 37NR <i>p</i> = 0.211	IHC	[174]
BRCA2	25	Neo-adjuvant locally advanced, recurrent	Docetaxel	Low: 5R vs 0NR High: 5R vs 15NR <i>p</i> = 0.0022	RT-PCR	[119]
ER	69	Primary or recurrent	Docetaxel	ER ⁺ : 18R vs 14NR ER ⁻ : 21R vs 16NR <i>P</i> = 0.862	IHC	[104]
	45	Neo-adjuvant primary	Docetaxel	ER ⁺ : 23R vs 9NRER ⁻ : 8R vs 5NR <i>P</i> = 0.502	IHC	[175]
	51	Neo-adjuvant primary	Docetaxel	ER ⁺ : 30R vs 3NR ER ⁻ : 9R vs 9NR <i>P</i> = 0.004	IHC	[176]
	100	Primary	Docetaxel	ER ⁺ : 4R vs 48NR ER ⁻ : 16R vs 32NR <i>P</i> = 0.002	IHC Response by pCR (pathological complete response)	[173]
GSTP1	62	Primary	Taxane	GST ⁺ : 0.31 ± 0.09 GST ⁻ : 0.73 ± 0.04 mean reduction rate <i>p</i> < 0.001	IHC Docetaxel significant Paclitaxel was NS	[159]
HER2	126	MBC	Taxane	pHer-2 ⁺ 1,2,3: 7CB vs 5PD pHer-2 ⁻ : 94CB vs 20PD <i>p</i> = 0.046	IHC of p1248Her-2	[177]
	66	MBC	Docetaxel	Her-2 ⁺ : 16R vs 14NR Her-2 ⁻ : 19R vs 17NR <i>P</i> = 0.5	IHC	[178]
	46	Neo-adjuvant primary	Docetaxel	Her-2 ⁺ : 10R vs 5NR Her-2 ⁻ : 21R vs 10NR <i>P</i> = 0.942	IHC	[175]
	62	Primary or recurrent	Docetaxel	Her-2 ⁺ : 9R vs 4NR Her-2 ⁻ : 24R vs 25NR <i>P</i> = 0.193	IHC	[104]
	67	Neo-adjuvant primary	Taxane	Her-2 ^{Amp} : 3R vs 16NRHer-2 ^{WT} : 5R vs 43NR <i>P</i> = 0.68	FISHResponse by pCR (pathological complete response)	[179]
	37	MBC	Docetaxel	Her-2 ⁺ : 6R vs 3NR Her-2 ⁻ : 9R vs 18NR <i>P</i> = 0.046	IHC	[180]
	100	Primary	Docetaxel	Her-2 ⁺ : 7R vs 19NR Her-2 ⁻ : 13R vs 61NR <i>P</i> = 0.305	IHC Response by pCR (pathological complete response)	[173]
	71	MBC	Docetaxel	Her-2 ^{Amp} : 14R vs 21NRHer-2 ^{WT} : 20R vs 50NR		[181]
HER2 ECD	35	MBC	Paclitaxel	HER2 ECD ⁺ : 40.9% HER2 ECD ⁻ : 38.5% <i>P</i> = 0.4	ELISA	[182]
Ki67	100	Primary	Docetaxel	Ki67 ⁺ : 17R vs 52NRKi67 ⁻ : 3R vs 28NR <i>P</i> = 0.108	IHC Response by pCR (pathological complete response)	[173]
p53	50	Primary or recurrent	Docetaxel	p53 ^{Mut} : 7R vs 9NR p53 ^{WT} : 21R vs 13NR NS	39/136 additional tumors p53 ^{Mut}	[183]
	63	Neo-adjuvant primary	Docetaxel	p53 high: 10R vs 22NR p53 low: 5R vs 24NR NS	IHC	[172]
	64	Primary or recurrent	Docetaxel	p53 ⁺ : 16R vs 13NR p53 ⁻ : 18R vs 15NR <i>p</i> = 0.961	IHC	[104]

Table 3 (continued)

Factor	Patient No	Patient Type	Treatment	Response	Comment	References
	114	Primary	paclitaxel	P53 ^{Mut} : 14R vs 11NR P53 ^{WT} : 36R vs 45NR $p = 0.1487$	Whole gene sequencing	[184]
Pgp	63	Primary or recurrent	Docetaxel	Pgp+: 14R vs 12NR Pgp-: 20R vs 17NR $P = 0.987$	IHC	[104]
Tau	41	MBC	Docetaxel	Tau ⁺ : 19R vs 13NR Tau ⁻ : 2R vs 2NR $P = 0.99$	IHC	[133]
	92	Primary	Docetaxel	Tau ⁺ : 8R vs 5NR Tau ⁻ : 9R vs 40NR $P = 0.977$	IHC Response by pCR (pathological complete response)	[173]
Thioredoxin	63	Neo-adjuvant primary	Docetaxel	Low: 15R vs 34NR High: 0R vs 14NR $p = 0.018$	IHC	[172]
	63	Primary or recurrent	Docetaxel	Thioredoxin ⁺ : 3R vs 11NR Thioredoxin ⁻ : 31R vs 18NR $P = 0.018$	IHC	[104]
β-tubulin isoforms	39	Locally advanced, recurrent	Docetaxel	Low β-I: 12R vs 7NR High β-I: 6R vs 14NR $p < 0.05$ Low β-III: 13R vs 6NR High β-III: 5R vs 15NR $p < 0.01$	RT-PCR	[135]
	41	MBC	Docetaxel	β-II ⁺ : 7R vs 11NR β-II ⁻ : 11R vs 3NR $P = 0.04$ β-III ⁺ : 11R vs 9NR β-III ⁻ : 7R vs 4NR $P = 0.72$ β-IV ⁺ : 19R vs 12NR β-IV ⁻ : 3R vs 4NR $P = 0.43$	IHC	[133]
	56	Primary or recurrent	Docetaxel	β-III ⁺ : 4R vs 10NR β-III ⁻ : 25R vs 17NR $P = 0.05$	IHC	[104]
	23	MBC	Paclitaxel	β-I WT: 11R vs 9NR β-I Mutation: 1R vs 3NR	β-I Mutational analysis (DHPLC-Seq)	[185]

MBC, metastatic breast cancer; R, response (generally CR + PR + SD); CB, clinical benefit; PD, progressive disease; NR, no response; IHC, immunohistochemistry; RT-PCR, reverse transcriptase polymerase chain reaction; NS, not significant; p53^{Mut}, mutation present; p53^{WT}, wild type p53; p53⁺, IHC positive according to criteria; p53⁻, IHC negative according to criteria; RT-PCR, reverse transcriptase polymerase chain reaction; ELISA, enzyme linked immunosorbent assay; FISH, fluorescent in situ hybridization; ECD, extracellular domain.

Note: only studies in which a taxane (single agent) was administered in chemotherapy naïve patients have been included; including studies where >10% of the inclusive population were not naïve.

codon 217 and another mutation, L215I resulting in enhanced binding of paclitaxel.⁹⁶ Mutations have also been reported in α-tubulin.⁷²

p53

The involvement of p53 in taxane resistance is very complex considering that wild type p53 leads to cell cycle arrest in the presence of DNA damage allowing for DNA repair. In the case of mutated p53 (mP53) it was expected that cells would be sensitive to DNA damaging agents, however, this is not always the case. Mutant p53 disables the apoptotic machinery often resulting in resistance to various drugs.^{100–103} However, to complicate the issue further it appears from a variety of sources that mP53 does not lead to paclitaxel or docetaxel resistance.¹⁰⁴ There are several lines of thought on this matter derived from both *in vitro* and *in vivo* data. In p53 KO (knock-out) mice increased sensitivity to paclitaxel is observed.¹⁰⁵ This is also seen in a number of p53 inactive cell lines⁹¹; however, the data are so varied and diverse there appears to be little if any consensus.

Indeed, there is no consensus on the range of exons analyzed with some authors analyzing exons 5–8 and others 4–9. In addition to this few studies incorporate analysis by more than one technique, they either use IHC (often with heterogeneous reporting criteria, and a range of different monoclonal antibodies) or perform mutational

analysis. Such differences in analytical techniques and heterogeneous assays continue to contribute to a lack of consensus being formed with regard to any clinical significance of p53, whether that be on account of overexpression or as a result of its mutational status.

Apoptosis

Considerable interest has been placed on deciphering the events associated with taxane related sensitivity and resistance through the study of apoptosis. Early studies indicated that over-expression of Bcl-2 and Bcl-xL contributed to taxane resistance. Additional studies led to the suggestion that there may be a threshold at which specific genes confer resistance.¹⁰⁶ There are other studies that indicate over-expression of pro-apoptotic genes are associated with paclitaxel sensitivity,¹⁰⁷ and yet others that show no correlation of Bcl-2 levels and response.¹⁰⁴

As most of the anti-apoptotic genes (*IAP*, *TRAF*, *Bcl-2*, *Bcl-xL*) are under the transcriptional control of NFκB this has also been studied. In fact it appears from a few studies that NFκB is constitutively activated in many breast cancers and that inhibition of NFκB may sensitize cells to taxanes.^{108–110} In addition, it has been shown that approximately half of breast cancers have increased levels of Akt, which appears to activate Bcl-2 and also increase the activation of NFκB.^{111–113} Akt is also activated by HER-2 signaling and is implicated in chemoresistance mechanisms of taxanes.^{114,115}

Cell cycle

The spindle assembly checkpoint (SAC) appears critical for taxane mediated cell death. Various mechanisms involved in this checkpoint appear to influence and be influenced by taxanes, and defects in the SAC correlate with resistance. (Fig. 1)

Various data sets seem to support this strong interaction:

- (1) Upon activation of the SAC both Mad2 and BubR1 interact with Cdc2 inhibiting its ability to activate APC.⁵⁰ Destruction of cyclin B and other regulators of mitosis by APC are responsible for proper metaphase-anaphase transition and mitotic exit. MAPs including Mad2 and BubR1 are thought to regulate SAC preventing anaphase until chromosomes are attached to bipolar spindles.¹¹⁶ In the presence of spindle inhibitors cyclin B degradation is inhibited, cells arrest at pro-metaphase and maintain constitutive Cdk1 activity (destruction of cyclin B inactivates Cdk1).⁵⁰
- (2) When Mad2 levels are low SAC is non-functional. Sensitivity is restored with re-establishment of Mad2 levels.⁵⁰
- (3) Over-expression of cyclins E and A have been associated with adverse outcomes. These cyclins are important mediators of G₁–S phase transition and subsequent S–G₂ phase transition. Cyclin A appears to be the more important as it is directly involved in regulating Cdk1 (cdc2) activity as activated Cdk1 is required for cells to enter mitosis and for SAC functionality, both key requirements of taxane sensitivity.¹¹⁷
- (4) BRCA1 is also implicated in SAC control. *BubR1* transcription is regulated by BRCA1, and also to some extent by p53. BRCA1 is a co-activator of p53 and positively regulates Mad2, thus inhibiting APC activity. Therefore, in BRCA1 deficient cells there is a premature onset of anaphase activated by APC by ubiquitination and degradation of cyclin B and subsequent activation of Cdk1,¹¹⁸ this being linked to paclitaxel resistance. Similarly there is data showing that decreased levels of BRCA2 correlate with better responses to docetaxel.¹¹⁹ (Table 3)
- (5) Stathmin, a microtubule regulator, destabilizes microtubules by two mechanisms, catalytic promotion and tubulin sequestration. It is active in G₂/M transition where it is inactivated by Cdk1 allowing for M phase entry. Stathmin over-expression has been correlated with resistance to taxanes.¹²⁰ More recently in a two-dimensional gel electrophoresis and MALDI-TOF peptide mass fingerprinting study stathmin was profiled as one of 9 proteins differentially expressed in paclitaxel resistant MCF-7 cells.¹²¹
- (6) HER2 over-expression inhibits taxol induced apoptosis by transcriptionally up-regulating p21^{cip1} which associates with p34^{Cdc2} inhibiting taxol mediated p34 activation delaying cells from entering G₂/M and thereby inhibiting apoptosis.¹²² HER2 may also directly phosphorylate Cdc2 leading to resistance. There is also evidence that HER2 positive tumors have low levels of Cdk1 resulting in delayed mitosis and paclitaxel resistance.¹²³ HER2 has also been shown to promote G₁/S progression and tumor cell proliferation by reducing p27^{Kip1} stability and reducing p27-cdc2 complexes. Stimulation through HER2 and other RTKs has also been shown to result in increased levels of Pgp, without affecting transcription, through the MEK–ERK–RSK pathway as inhibitors to this pathway decrease Pgp mediated resistance to paclitaxel.¹²³

Thus it appears that the microtubule composition of the mitotic spindle, the dynamics of microtubule assembly and the associated anaphase–metaphase block induced by anti-microtubule agents highlight the importance of the transition into M phase in deter-

mining their sensitivity and that deficits in any of the multitude of proteins that regulate the SAC would be sufficient for conferring resistance to taxanes.

Gene signatures/metagenes

Several groups have investigated high throughput screening of thousands of genes as a method to identify patterns of expression of single genes or gene combinations (gene signatures) that correlate with outcome to given therapies. A first study to investigate taxane related outcomes of response in breast cancer was reported by Chang, et al.¹²⁴ They identified 92 genes that correlated with docetaxel response of primary breast cancer in the neo-adjuvant setting. Their RNA profile from 24 patients included higher expression of genes involved in cell cycle, cell adhesion, protein modification, transcription and apoptosis; while resistant tumors showed increased expression of some transcriptional and signal transduction genes. The 92 gene predictor had positive and negative predictive values of 92% and 83%, respectively.¹²⁴

Utilizing another technique, adaptor-tagged competitive (ATAC)-polymerase chain reaction (PCR), Iwao-Koizumi et al., measured the expression of 2453 genes in a series of 70 (44 learning set, 26 validation set) primary or locally recurrent breast cancers receiving docetaxel.¹²⁵ They identified an algorithm consisting of 85 genes that predicted clinical response to docetaxel with positive and negative predictive values of 73.3% and 90.9%, respectively. Non-responders were characterized as having elevated expression of genes controlling cellular redox, thioredoxin, glutathione-S-transferase and peroxiredoxin.¹²⁵

The utility of gene expression signatures based predictive algorithms will advance as they allow, if standardized, potentially improved positive and negative predictive ability over single gene predictors. Similar algorithms of metagenes may also enter into clinical development as our knowledge base increases per predictive marker/signature.

Clinically relevant prediction of taxane resistance

While the majority of data presented relates to *in vitro* experimentation, limited hypothesis of taxane resistance have been investigated *in vivo*. Therefore, insufficient *in vivo* data exist to gain a clear picture of numerous hypotheses that have been generated for molecules of predictive/prognostic significance in breast cancer. To date studies have failed to indicate any particular mechanism or marker as immediately clinically relevant with respect to offering insights into patient stratification.

P-glycoprotein (Pgp)

Increased expression in breast cancers and in other cancer types has generally been correlated with MDR.⁶⁸ However, much of the data is conflicting. Some of the problems in assessing the predictive nature of Pgp may relate to the variety of methods used and lack of standardization of cut offs for quantitation, clinical endpoints measured and study/patient heterogeneity that exists across all studies. Furthermore, one of the main antibodies used for its detection has been reported to cross react with HER2 and also the heavy chain of myosin leading to distinct difficulties in interpretation.¹²⁶

Putting this aside Pgp expression (depending on the method of analysis and thresholds used) shows a broad range of expression. From 0–30% in newly diagnosed breast cancer rising to over 70% in many cases of relapsed breast cancer.¹²⁷ Indeed in a meta-analysis of MDR1/Pgp expression in breast cancers Trock et al., showed that approximately 40% of breast tumors expressed Pgp at RNA level or protein assessed by IHC, and that in tumors

analyzed post chemotherapy the incidence of Pgp positivity increased.¹²⁸ Furthermore, patients with Pgp positive tumors were three times less likely to achieve an objective response compared to those with Pgp negative tumors. Nevertheless, a word of caution needs to be raised here as very few of the included studies represent patients treated with single agent taxane, there was great heterogeneity between study populations and the methods of expression were also inconsistent.

In one study examining the relationship between YB-1, a Y-box binding protein that targets the Pgp promoter, YB-1 localization and Pgp expression indicated that translocation of YB-1 from the cytoplasm to the nucleus was associated with increased expression of Pgp in breast cancers.¹²⁹ The authors claimed that nuclear localization of YB-1 was significantly correlated with resistance to paclitaxel. Their data confirmed previous reports indicating that YB-1 is important in drug sensitivity of cancers via increasing Pgp levels, offering another mechanism for increased Pgp in chemotherapy treated breast cancers.

The development of alternative technologies for measuring P-gp expression real-time *in vivo* include Technetium-^{99m}-Sestamib radio-imaging, however, it has not as yet offered clinical prediction of response to paclitaxel.⁹¹ Additional efflux pumps other than Pgp have also been investigated, again with limited predictive ability.

β -Tubulin

With regards to the possible alterations in tubulin levels or the dynamics of tubulin polymerization being related to taxane resistance in the clinical setting remains a more difficult challenge than demonstrated *in vitro*. Studies investigating both alterations in β -tubulin isoform expression levels and somatic mutations have proved to be technically demanding and have generated conflicting results in all tumor types investigated, one reason for these differences being the existence of nine β -tubulin pseudogenes and all share substantial sequence homology with the functional gene.^{130–132}

As indicated in Table 3 only few studies have investigated β -tubulin isoforms in breast cancer treated with single agent taxanes. Two of the three studies used IHC, one RT-PCR and one examined the mutational status. In the study by Noguchi¹⁰⁴ β -III levels reached borderline statistical significance, however, this is not shared in the study by Bernard-Marty et al.¹³³ Their study indicated the significance of β -I and β -VI, β -I has also shown significance by RT-PCR.¹³⁴ Differences in the techniques utilized and the populations studied (MBC versus primary or recurrent versus locally advanced, recurrent) again limit the strength of any cumulative data related to the potential clinical utility of β -tubulin isoforms.

Tau

Tau is a stabilizer of microtubule assembly. One interesting feature of Tau is that over-expression is linked to ER expression. The gene for Tau also contains an ER responsive element, thus questioning the predictive nature of tau expression with endocrine sensitivity. At a functional level tau binds to the same microtubule binding pocket that taxanes bind, indicating that low tau expression should correlate with taxane sensitivity, while high tau expression may indicate increased benefit from endocrine therapy.¹³⁵ Limited data from clinical data sets exist regarding tau, see Table 3.

BRCA

Recently “BRCAness” has come to the forefront with the correlation of triple negative (HER-2, ER and PR negative tumors) breast

cancers (TNBC) and hereditary breast cancers harboring mutations in *BRCA1* or *BRCA2*.¹³⁶ Following on from this several studies have indicated that chemo-sensitivity of TNBC may be higher to non-taxane and non-anthracycline containing regimens,^{94–96} although there is contradictory data.^{137–139} At this early point in time it would be advisable to await additional and better designed analyses of appropriate studies before conclusions are made regarding the utility of *BRCA* status for guiding treatment with taxanes. Other groups are also investigating meta-genes for the identification of signatures that may predict response to taxanes in TNBC.¹⁴⁰

HER2

In vitro experiments of systems with HER2 over-expression indicate that down-regulation of HER2 using neutralizing antibodies (including Herceptin) can reverse the resistance observed with taxanes. Other reports indicate that HER2 may confer resistance independent of MDR-1. Herceptin demonstrates tumor inhibitory and chemo-sensitizing effects with paclitaxel,⁵⁵ and docetaxel.^{104,141} The mechanism of resistance to taxanes in HER2 over-expressing tumors is unknown, however, it is suggested that HER2 over-expression induces resistance by increasing p21 expression leading to CDK1 inhibition, that results in blockade of taxane mediated apoptosis.¹⁴² (see Fig. 1)

Several issues of HER2 status could lead to misinterpretation of the published data sets. It has long been recognized that *HER2* amplification is associated with co-amplification of the *Topoisomerase IIa* gene in 40–50% of cases.^{143,144} This could be a factor that biases results obtained from both standard IHC but also from FISH based analyses, and likewise any chemosensitivity/ resistance on behalf of the *TOPOII* gene would also need to be investigated independently from HER2 expression or gene amplification.^{145,146}

As indicated in Table 3, although there is speculation that HER2 overexpression, upregulation or gene gain may be a potential biomarker of taxane resistance; the few studies that have assessed this hypothesis do not demonstrate a consistent trend. Two of the eight studies demonstrated a difference in MBC patients and one showed border-line significance. Unfortunately the number of studies is limited, the populations are relatively small and no homogeneous stratification by technique was employed. Today the establishment of the addition of Herceptin to taxanes in the adjuvant setting, sequential often utilized in the European Union, and concomitant in the USA, will likely make such an analysis of a prospective study unlikely.

Other factors

While there are a number of other predictive factors for response to taxanes it is impossible to list all of those reported in detail.^{91,104} The principle mechanisms are highlighted in Table 3 along with some of the more interesting of the novel mechanisms.

Circumvention

Resistance to taxanes parallels that of resistance to chemotherapy, occurring prior to exposure (*de-novo*, or innate), or as a result of exposure (acquired resistance). In addition there is what is termed cross-resistance or multi-drug resistance, where individuals exposed to one specific compound develop resistance to multiple structurally unrelated compounds. In the case of taxanes all three mechanisms have been reported. Once these forms of taxane resistance become evident few subsequent treatment options are available for breast cancer patients. The majority of such patients are subsequently treated with capecitabine, gemcitabine and vinorelbine, however, the response rates remain low.¹⁴⁷

Several strategies have been used, with limited success with respect to circumventing taxane resistance. The epithiliones, a family of naturally occurring cytotoxic microtubule inhibitors, represent the most advanced agents for taxane resistant or refractory patients. Epithiliones are structurally different from paclitaxel and docetaxel having a different mechanism of action. They stabilize microtubules, suppress their dynamics, induce mitotic arrest in G₂/M resulting in apoptosis similar to that of the effect of taxanes.¹⁴⁸ Although they bind to similar sites as the taxanes, they have a unique and independent molecular interaction that is partly to explain their resistance to the classical multidrug resistance mechanisms.¹⁴⁹

A number of different anti-tubulin agents have progressed into phase I clinical trials or are entering late phase III analysis including: epithilione B (patupilone); epithilione D; hailchondrin B; taxane analog DJ-927 and ixabepilone.¹⁵⁰ Ixabepilone is the most developed of these having demonstrated low resistance (*in vivo* and *in vitro*) to various drug resistance mechanisms that affect the taxanes including: MDR overexpression [81 of 1], β -tubulin mutations, β -III isotype overexpression. It has also been shown that ixabepilone weekly induces Pgp expression; and retains the ability to bind to β -III microtubules.¹⁵¹ Following clinical analysis in several phase II and phase III studies it was the first epithilione to be approved by the FDA. It is indicated in combination with capecitabine for the treatment of MBC, or locally advanced breast cancer resistant or refractory to treatment with an anthracycline and a taxane, or as monotherapy for the treatment of MBC or locally advanced breast cancer resistant or refractory to anthracyclines, taxanes and capecitabine.^{150,152–156} The search for additional agents that do not share the resistance mechanisms of the taxanes is a continuing process.

Discussion/conclusions

Some of the most obvious limitations to understanding the clinical relevance of any of the aforementioned mechanisms of resistance relate to the lack of universally accepted guidelines for analytical and/or clinical validation, differences in methods of tissue collection preparation and storage, different target assays being utilized and the associated problems of sensitivity and specificity of currently used immunohistochemical analyses.^{157,158} Even in the most comprehensive meta-analysis to date of Pgp in breast cancer, study heterogeneity, sample and analytical heterogeneity significantly limited the extraction of reliable data.⁶⁸

As indicated from the data sets presented for taxane treated chemo-naïve breast cancer patients the cumulative best evidence for any suggested biomarker is limited but to a few studies. It is obvious that there are severe weaknesses such as retrospective analysis, sample size and lack of clinical information beyond that of response. For these reasons the real predictive value of any of the candidate biomarkers remains ill-defined. Furthermore, the utility of individual biomarkers remains a limited approach as many of the agents do not have only one target, and thus multiple gene models or signatures may be more informative. Unraveling potential differences between for example Paclitaxel and Docetaxel are further restricted. Most of the data tabulated in this review pertains to docetaxel treated populations (Table 3). Only one study reported on differences between the two groups, with a significant difference in response found between high and low IHC expression levels of GSTP1 for docetaxel alone.¹⁵⁹ No supportive data exists for GSTP1 and limitations in the entire field of Taxane Resistance Biomarkers does not support the idea that the drugs resistance profiles are different.

There are also several specific issues related to some of the individual candidate biomarkers and their rationale in light of our

understanding of taxane function. If we consider that the taxanes function by stabilizing microtubules leading to cell cycle arrest at G₂/M (Fig. 1) and subsequent apoptosis, biomarkers of resistance should have some functional interaction in this process. Taxanes attach to the β -subunit of tubulin and therein do not directly cause DNA-damage. With this in mind there is little supposed benefit from the study of p53 status as docetaxel induced cell cycle arrest occurs in a late phase of G₂/M. Mutational analysis of p53 is also thwart with errors simply as we do not know exactly what each mutant variant does. Some generate stable aberrant protein, and others generate no protein at all.¹⁶⁰ Furthermore, some mutation positive cases may generate mutant p53 that is undetectable by a given antibody clone used for IHC.

One trend that may be possible to discuss is that of the association between taxane efficacy and the proliferation index of the tumor. Indeed most chemotherapeutic agents work better in cancers with a high proliferative index.^{161–163} This high proliferative index characterized by rapidly growing tumors may therein correlate with some of the candidate biomarkers indicated in Table 3. Unfortunately there is little collaborative data associated with this possibility as only a few studies have examined a routine set of markers (e.g. HER2, ER, Ki67) and grade is not one of the factors stratified in the studies reviewed herein.

It appears from the limited number of published studies that there are no valid practical biomarkers that could predict resistance to the taxanes. There remains an immediate clinical requirement for a biomarker of taxane resistance. The community will need to come together in addressing several consanguineous candidate biomarkers, however, there is also a need for the community to understand that eligible data sets will only be those that have received a taxane (or possibly a taxane containing regimen) in predefined chemotherapy naïve patient populations. Some examples of possible data sets that could be analyzed retrospectively include: ECOG 2100,¹⁶⁴ E1193¹⁶⁵ and others.^{166–171} However, it is only by the incorporation of such biomarker analyses into prospectively designed studies that clinical practice will alter; such studies are awaited.

Conflict of Interest

Consultant or Advisory role: Dr. S. Murray, Merck KGaA, Darmstadt, Germany. Merck distribute the MoAb Cetuximab (Erbix[®]). Dr. S. Murray and Dr. P. Kosmidis, AstraZeneca, Macclesfield, United Kingdom. AstraZeneca are proprietors of gefitinib (Iressa[®]). Dr. S. Murray, Amgen Thousand OaksCa, USA. Amgen distribute the MoAb Panitumumab (Vectibix[®]). No other author has a conflict of interest.

References

1. Jemal A, Bray F, Center MM, Ferlay J, Ward E, Forman D. Global cancer statistics. *CA Cancer J Clin* 2011;**61**(2):69–90.
2. <http://seer.cancer.gov/statfacts/html/breast.html>.
3. King KM, Lupichuk S, Baig L, Webster M, Basi S, Whyte D, et al. Optimal use of taxanes in metastatic breast cancer. *Curr Oncol* 2009;**16**(3):8–20.
4. Chan S, Friedrichs K, Noel D, Pinter T, Van Belle S, Vorobiof D, et al. Prospective randomized trial of docetaxel versus doxorubicin in patients with metastatic breast cancer. *J Clin Oncol* 1999;**17**(8):2341–54.
5. Evans TR, Yellowlees A, Foster E, Earl H, Cameron DA, Hutcheon AW, et al. Phase III randomized trial of doxorubicin and docetaxel versus doxorubicin and cyclophosphamide as primary medical therapy in women with breast cancer: an anglo-celtic cooperative oncology group study. *J Clin Oncol* 2005;**23**(13):2988–95.
6. Nabholz JM, Senn HJ, Bezwoda WR, Melnychuk D, Deschenes L, Douma J, et al. Prospective randomized trial of docetaxel versus mitomycin plus vinblastine in patients with metastatic breast cancer progressing despite previous anthracycline-containing chemotherapy. 304 Study Group. *J Clin Oncol* 1999;**17**(5):1413–24.
7. Jassem J, Pienkowski T, Pluzanska A, Jelic S, Gorbunova V, Mrsic-Krmpotic Z, et al. Doxorubicin and paclitaxel versus fluorouracil, doxorubicin, and

- cyclophosphamide as first-line therapy for women with metastatic breast cancer: final results of a randomized phase III multicenter trial. *J Clin Oncol* 2001;**19**(6):1707–15.
8. Bedard PL, Di Leo A, Piccart-Gebhart MJ. Taxanes: optimizing adjuvant chemotherapy for early-stage breast cancer. *Nat Rev Clin Oncol* 2010;**7**(1):22–36.
 9. Palmieri C, Krell J, James CR, Harper-Wynne C, Misra V, Cleator S, et al. Rechallenge with anthracyclines and taxanes in metastatic breast cancer. *Nat Rev Clin Oncol* 2010;**7**(10):561–74.
 10. Cardoso F, Senkus-Konefka E, Fallowfield L, Costa A, Castiglione M. Locally recurrent or metastatic breast cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2010;**21**(Suppl. 5):v15–19.
 11. Gonzalez-Angulo AM, Morales-Vasquez F, Hortobagyi GN. Overview of resistance to systemic therapy in patients with breast cancer. *Adv Exp Med Biol* 2007;**608**:1–22.
 12. Pohlmann PR, Mayer IA, Mernaugh R. Resistance to trastuzumab in breast cancer. *Clin Cancer Res* 2009;**15**(24):7479–91.
 13. Zhao Y, Liu H, Liu Z, Ding Y, Ledoux SP, Wilson GL, et al. Overcoming trastuzumab resistance in breast cancer by targeting dysregulated glucose metabolism. *Cancer Res* 2011;**71**(13):4585–97.
 14. Dalton WS. Mechanisms of drug resistance in breast cancer. *Semin Oncol* 1990;**17**(4 Suppl. 7):37–9.
 15. Rottenberg S, Nygren AO, Pajic M, van Leeuwen FW, van der Heijden I, van de Wetering K, et al. Selective induction of chemotherapy resistance of mammary tumors in a conditional mouse model for hereditary breast cancer. *Proc Natl Acad Sci U S A* 2007;**104**(29):12117–22.
 16. O'Brien C, Cavet G, Pandita A, Hu X, Haydu L, Mohan S, et al. Functional genomics identifies ABCC3 as a mediator of taxane resistance in HER2-amplified breast cancer. *Cancer Res* 2008;**68**(13):5380–9.
 17. Brooks TA, Minderman H, O'Loughlin KL, Pera P, Ojima I, Baer MR, et al. Taxane-based reversal agents modulate drug resistance mediated by P-glycoprotein, multidrug resistance protein, and breast cancer resistance protein. *Mol Cancer Ther* 2003;**2**(11):1195–205.
 18. Raguz S, Yague E. Resistance to chemotherapy: new treatments and novel insights into an old problem. *Br J Cancer* 2008;**99**(3):387–91.
 19. Ahmed-Belkacem A, Pozza A, Munoz-Martinez F, Bates SE, Castanys S, Gamarro F, et al. Flavonoid structure-activity studies identify 6-prenylchrysin and tectochrysin as potent and specific inhibitors of breast cancer resistance protein ABCG2. *Cancer Res* 2005;**65**(11):4852–60.
 20. Kuhnle M, Egger M, Muller C, Mahringer A, Bernhardt G, Fricker G, et al. Potent and selective inhibitors of breast cancer resistance protein (ABCG2) derived from the p-glycoprotein (ABCB1) modulator tariquidar. *J Med Chem* 2009;**52**(4):1190–7.
 21. Yoshida N, Takada T, Yamamura Y, Adachi I, Suzuki H, Kawakami J. Inhibitory effects of terpenoids on multidrug resistance-associated protein 2- and breast cancer resistance protein-mediated transport. *Drug Metab Dispos* 2008;**36**(7):1206–11.
 22. Bollag DM, McQueney PA, Zhu J, Hensens O, Koupal L, Liesch J, et al. Etoposides, a new class of microtubule-stabilizing agents with a taxol-like mechanism of action. *Cancer Res* 1995;**55**(11):2325–33.
 23. Rivera E, Lee J, Davies A. Clinical development of ixabepilone and other epothilones in patients with advanced solid tumors. *Oncologist* 2008;**13**(12):1207–23.
 24. Farmer P, Bonnefoi H, Anderle P, Cameron D, Wirapati P, Becette V, et al. A stroma-related gene signature predicts resistance to neoadjuvant chemotherapy in breast cancer. *Nat Med* 2009;**15**(1):68–74.
 25. Harris LN, You F, Schnitt SJ, Witkiewicz A, Lu X, Sgroi D, et al. Predictors of resistance to preoperative trastuzumab and vinorelbine for HER2-positive early breast cancer. *Clin Cancer Res* 2007;**13**(4):1198–207.
 26. Pandolfi PP. Breast cancer—loss of PTEN predicts resistance to treatment. *N Engl J Med* 2004;**351**(22):2337–8.
 27. Lee CH, Macgregor PF. Using microarrays to predict resistance to chemotherapy in cancer patients. *Pharmacogenomics* 2004;**5**(6):611–25.
 28. Linardou H, Briassoulis E, Dahabreh IJ, Mountzios G, Papadimitriou C, Papadopoulos S, et al. All about KRAS for clinical oncology practice: gene profile, clinical implications and laboratory recommendations for somatic mutational testing in colorectal cancer. *Cancer Treat Rev* 2011;**37**(3):221–33.
 29. Joerger M, Huitema AD, Richel DJ, Dittich C, Pavlidis N, Briassoulis E, et al. Population pharmacokinetics and pharmacodynamics of paclitaxel and carboplatin in ovarian cancer patients: a study by the European organization for research and treatment of cancer-pharmacology and molecular mechanisms group and new drug development group. *Clin Cancer Res* 2007;**13**(21):6410–8.
 30. Thomas C, Gustafsson JA. A CUE hints at tumor resistance. *Nat Med* 2011;**17**(6):658–60.
 31. Rowinsky EK, Donehower RC. Paclitaxel (taxol). *N Engl J Med* 1995;**332**(15):1004–14.
 32. Thigpen JT, Blessing JA, Ball H, Hummel SJ, Barrett RJ. Phase II trial of paclitaxel in patients with progressive ovarian carcinoma after platinum-based chemotherapy: a Gynecologic Oncology Group study. *J Clin Oncol* 1994;**12**(9):1748–53.
 33. Einzig AI. Review of phase II trials of Taxol (paclitaxel) in patients with advanced ovarian cancer. *Ann Oncol* 1994;**5**(Suppl. 6):S29–32.
 34. Fountzilias G, Athanassiades A, Giannakakis T, Bafaloukos D, Karakousis K, Dombros N, et al. A phase II study of paclitaxel in advanced breast cancer resistant to anthracyclines. *Eur J Cancer* 1996;**32A**(1):47–51.
 35. Wilson WH, Berg SL, Bryant G, Wittes RE, Bates S, Fojo A, et al. Paclitaxel in doxorubicin-refractory or mitoxantrone-refractory breast cancer: a phase I/II trial of 96-hour infusion. *J Clin Oncol* 1994;**12**(8):1621–9.
 36. Millward MJ, Bishop JF, Friedlander M, Levi JA, Goldstein D, Oliver IN, et al. Phase II trial of a 3-hour infusion of paclitaxel in previously untreated patients with advanced non-small-cell lung cancer. *J Clin Oncol* 1996;**14**(1):142–8.
 37. Gatzemeier U, Heckmayr M, Neuhauss R, Schluter I, Pawel JV, Wagner H, et al. Phase II study with paclitaxel for the treatment of advanced inoperable non-small cell lung cancer. *Lung cancer* 1995;**12**(Suppl. 2):S101–106.
 38. Papamichael D, Gallagher CJ, Oliver RT, Johnson PW, Waxman J. Phase II study of paclitaxel in pretreated patients with locally advanced/metastatic cancer of the bladder and ureter. *Br J Cancer* 1997;**75**(4):606–7.
 39. Ringel I, Horwitz SB. Studies with RP 56976 (taxotere): a semisynthetic analogue of taxol. *J Natl Cancer Inst* 1991;**83**(4):288–91.
 40. Gligrov J, Lotz JP. Preclinical pharmacology of the taxanes: implications of the differences. *Oncologist* 2004;**9**(Suppl. 2):3–8.
 41. Parness J, Horwitz SB. Taxol binds to polymerized tubulin in vitro. *J Cell Biol* 1981;**91**(2 Pt 1):479–87.
 42. Rao S, Orr GA, Chaudhary AG, Kingston DG, Horwitz SB. Characterization of the taxol binding site on the microtubule. 2-(m-Azidobenzoyl)taxol photolabels a peptide (amino acids 217–231) of beta-tubulin. *J Biol Chem* 1995;**270**(35):20235–8.
 43. Crossin KL, Carney DH. Microtubule stabilization by taxol inhibits initiation of DNA synthesis by thrombin and by epidermal growth factor. *Cell* 1981;**27**(2 Pt 3):341–50.
 44. Fabbri F, Carloni S, Brigliadori G, Zoli W, Lapalombella R, Marini M. Sequential events of apoptosis involving docetaxel, a microtubule-interfering agent: a cytometric study. *BMC Cell Biol* 2006;**7**:6.
 45. Ferlini C, Cicchillitti L, Raspaglio G, Bartollino S, Cimitan S, Bertucci C, et al. Paclitaxel directly binds to Bcl-2 and functionally mimics activity of Nur77. *Cancer Res* 2009;**69**(17):6906–14.
 46. Varbiro G, Veres B, Gallyas Jr F, Sumegi B. Direct effect of Taxol on free radical formation and mitochondrial permeability transition. *Free Radic Biol Med* 2001;**31**(4):548–58.
 47. Ibrado AM, Kim CN, Bhalla K. Temporal relationship of CDK1 activation and mitotic arrest to cytosolic accumulation of cytochrome C and caspase-3 activity during Taxol-induced apoptosis of human AML HL-60 cells. *Leukemia* 1998;**12**(12):1930–6.
 48. Ling YH, Consoli U, Tornos C, Andreeff M, Perez-Soler R. Accumulation of cyclin B1, activation of cyclin B1-dependent kinase and induction of programmed cell death in human epidermoid carcinoma KB cells treated with taxol. *Int J Cancer* 1998;**75**(6):925–32.
 49. Yuan J, Kramer A, Matthess Y, Yan R, Spankuch B, Gatie R, et al. Stable gene silencing of cyclin B1 in tumor cells increases susceptibility to taxol and leads to growth arrest in vivo. *Oncogene* 2006;**25**(12):1753–62.
 50. Sudo T, Nitta M, Saya H, Ueno NT. Dependence of paclitaxel sensitivity on a functional spindle assembly checkpoint. *Cancer Res* 2004;**64**(7):2502–8.
 51. Berchem GJ, Bosseler M, Mine N, Avalosse B. Nanomolar range docetaxel treatment sensitizes MCF-7 cells to chemotherapy induced apoptosis, induces G2M arrest and phosphorylates bcl-2. *Anticancer Res* 1999;**19**(1A):535–40.
 52. Wang S, Wang Z, Boise L, Dent P, Grant S. Loss of the bcl-2 phosphorylation loop domain increases resistance of human leukemia cells (U937) to paclitaxel-mediated mitochondrial dysfunction and apoptosis. *Biochem Biophys Res Commun* 1999;**259**(1):67–72.
 53. Jordan MA, Toso RJ, Thrower D, Wilson L. Mechanism of mitotic block and inhibition of cell proliferation by taxol at low concentrations. *Proc Natl Acad Sci U S A* 1993;**90**(20):9552–6.
 54. Rudner AD, Murray AW. The spindle assembly checkpoint. *Curr Opin Cell Biol* 1996;**8**(6):773–80.
 55. Long BH, Fairchild CR. Paclitaxel inhibits progression of mitotic cells to G1 phase by interference with spindle formation without affecting other microtubule functions during anaphase and telophase. *Cancer Res* 1994;**54**(16):4355–61.
 56. Wang J, Lou P, Lesniewski R, Henkin J. Paclitaxel at ultra low concentrations inhibits angiogenesis without affecting cellular microtubule assembly. *Anticancer Drugs* 2003;**14**(1):13–9.
 57. Wang TH, Wang HS, Soong YK. Paclitaxel-induced cell death: where the cell cycle and apoptosis come together. *Cancer* 2000;**88**(11):2619–28.
 58. Gottesman MM, Hrycyna CA, Schoenlein PV, Germann UA, Pastan I. Genetic analysis of the multidrug transporter. *Annu Rev Genet* 1995;**29**:607–49.
 59. Baldini N. Multidrug resistance—a multiplex phenomenon. *Nat Med* 1997;**3**(4):378–80.
 60. Lee AJ, Endesfelder D, Rowan AJ, Walther A, Birkbak NJ, Futreal PA, et al. Chromosomal instability confers intrinsic multidrug resistance. *Cancer Res* 2011;**71**(5):1858–70.
 61. Keshelava N, Zuo JJ, Chen P, Waidyaratne SN, Luna MC, Gomer CJ, et al. Loss of p53 function confers high-level multidrug resistance in neuroblastoma cell lines. *Cancer Res* 2001;**61**(16):6185–93.
 62. Marquardt D, McCrone S, Center MS. Mechanisms of multidrug resistance in HL60 cells: detection of resistance-associated proteins with antibodies against synthetic peptides that correspond to the deduced sequence of P-glycoprotein. *Cancer Res* 1990;**50**(5):1426–30.
 63. Keizer HG, Schuurhuis GJ, Broxterman HJ, Lankelma J, Schoonen WG, van Rijn J, et al. Correlation of multidrug resistance with decreased drug accumulation, altered subcellular drug distribution, and increased P-glycoprotein expression

- in cultured SW-1573 human lung tumor cells. *Cancer Res* 1989;**49**(11):2988–93.
64. Zhu W, Shan X, Wang T, Shu Y, Liu P. MiR-181b modulates multidrug resistance by targeting BCL2 in human cancer cell lines. *Int J Cancer* 2010;**127**(11):2520–9.
 65. Ling V, Charles F. Kettering Prize. P-glycoprotein and resistance to anticancer drugs. *Cancer* 1992;**69**(10):2603–9.
 66. Schinkel AH. The physiological function of drug-transporting P-glycoproteins. *Semin Cancer Biol* 1997;**8**(3):161–70.
 67. Cordon-Cardo C, O'Brien JP, Boccia J, Casals D, Bertino JR, Melamed MR. Expression of the multidrug resistance gene product (P-glycoprotein) in human normal and tumor tissues. *J Histochem Cytochem* 1990;**38**(9):1277–87.
 68. Trock BJ, Leonessa F, Clarke R. Multidrug resistance in breast cancer: a meta-analysis of MDR1/gp170 expression and its possible functional significance. *J Natl Cancer Inst* 1997;**89**(13):917–31.
 69. Beck WT, Grogan TM, Willman CL, Cordon-Cardo C, Parham DM, Kuttlesch JF, et al. Methods to detect P-glycoprotein-associated multidrug resistance in patients' tumors: consensus recommendations. *Cancer Res* 1996;**56**(13):3010–20.
 70. Dean M, Rzhetsky A, Allikmets R. The human ATP-binding cassette (ABC) transporter superfamily. *Genome Res* 2001;**11**(7):1156–66.
 71. Gottesman MM, Fojo T, Bates SE. Multidrug resistance in cancer: role of ATP-dependent transporters. *Nat Rev Cancer* 2002;**2**(1):48–58.
 72. Fojo AT, Menefee M. Microtubule targeting agents: basic mechanisms of multidrug resistance (MDR). *Semin Oncol* 2005;**32**(6 Suppl. 7):S3–8.
 73. Szakacs G, Annereau JP, Lababidi S, Shankavaram U, Arciello A, Bussey KJ, et al. Predicting drug sensitivity and resistance. profiling ABC transporter genes in cancer cells. *Cancer cell* 2004;**6**(2):129–37.
 74. Kruh GD, Belinsky MG. The MRP family of drug efflux pumps. *Oncogene* 2003;**22**(47):7537–52.
 75. Gandhi YA, Morris ME. Structure-activity relationships and quantitative structure-activity relationships for breast cancer resistance protein (ABCG2). *Aaps J* 2009;**11**(3):541–52.
 76. Mao Q, Unadkat JD. Role of the breast cancer resistance protein (ABCG2) in drug transport. *Aaps J* 2005;**7**(1):E118–133.
 77. Sanfilippo O, Ronchi E, De Marco C, Di Fronzo G, Silvestrini R. Expression of P-glycoprotein in breast cancer tissue and in vitro resistance to doxorubicin and vincristine. *Eur J Cancer* 1991;**27**(2):155–8.
 78. Clarke R, Leonessa F, Trock B. Multidrug resistance/P-glycoprotein and breast cancer: review and meta-analysis. *Semin Oncol* 2005;**32**(6 Suppl. 7):S9–15.
 79. Merkel DE, Fuqua SA, McGuire WL. P-glycoprotein in breast cancer. *Cancer Treat Res* 1989;**48**:97–105.
 80. Bell DR, Trent JM, Willard HF, Riordan JR, Ling V. Chromosomal location of human P-glycoprotein gene sequences. *Cancer Genet Cytogenet* 1987;**25**(1):141–8.
 81. Osborn M, Weber K. The display of microtubules in transformed cells. *Cell* 1977;**12**(3):561–71.
 82. Saito SY. Toxins affecting actin filaments and microtubules. *Prog Mol Subcell Biol* 2009;**46**:187–219.
 83. Wade RH. Microtubules: an overview. *Methods Mol Med* 2007;**137**:1–16.
 84. Kerssemakers JW, Munteanu EL, Laan L, Noetzel TL, Janson ME, Dogterom M. Assembly dynamics of microtubules at molecular resolution. *Nature* 2006;**442**(7103):709–12.
 85. Dumontet C, Sikic BI. Mechanisms of action of and resistance to antitubulin agents: microtubule dynamics, drug transport, and cell death. *J Clin Oncol* 1999;**17**(3):1061–70.
 86. Mitchison T, Kirschner M. Dynamic instability of microtubule growth. *Nature* 1984;**312**(5991):237–42.
 87. Dutcher SK. The tubulin fraternity: alpha to eta. *Curr Opin Cell Biol* 2001;**13**(1):49–54.
 88. Stearns T, Evans L, Kirschner M. Gamma-tubulin is a highly conserved component of the centrosome. *Cell* 1991;**65**(5):825–36.
 89. Verhey KJ, Gaertig J. The tubulin code. *Cell Cycle* 2007;**6**(17):2152–60.
 90. Snyder JP, Nettles JH, Cornett B, Downing KH, Nogales E. The binding conformation of Taxol in beta-tubulin: a model based on electron crystallographic density. *Proc Natl Acad Sci U S A* 2001;**98**(9):5312–6.
 91. Yusuf RZ, Duan Z, Lamendola DE, Penson RT, Seiden MV. Paclitaxel resistance. molecular mechanisms and pharmacologic manipulation. *Curr Cancer Drug Targets* 2003;**3**(1):1–19.
 92. Han EK, Gehrke L, Tahir SK, Credo RB, Cherian SP, Sham H, et al. Modulation of drug resistance by alpha-tubulin in paclitaxel-resistant human lung cancer cell lines. *Eur J Cancer* 2000;**36**(12):1565–71.
 93. Minotti AM, Barlow SB, Cabral F. Resistance to antimetabolic drugs in Chinese hamster ovary cells correlates with changes in the level of polymerized tubulin. *J Biol Chem* 1991;**266**(6):3987–94.
 94. Rouzier R, Rajan R, Wagner P, Hess KR, Gold DL, Stec J, et al. Microtubule-associated protein tau: a marker of paclitaxel sensitivity in breast cancer. *Proc Natl Acad Sci U S A* 2005;**102**(23):8315–20.
 95. Giannakakou P, Sackett DL, Kang YK, Zhan Z, Buters JT, Fojo T, et al. Paclitaxel-resistant human ovarian cancer cells have mutant beta-tubulins that exhibit impaired paclitaxel-driven polymerization. *J Biol Chem* 1997;**272**(27):17118–25.
 96. Kirschner MW. Microtubule assembly and nucleation. *Int Rev Cytol* 1978;**54**:1–71.
 97. Banerjee A, Roach MC, Trcka P, Luduena RF. Preparation of a monoclonal antibody specific for the class IV isotype of beta-tubulin. Purification and assembly of alpha beta II, alpha beta III, and alpha beta IV tubulin dimers from bovine brain. *J Biol Chem* 1992;**267**(8):5625–30.
 98. Derry WB, Wilson L, Khan IA, Luduena RF, Jordan MA. Taxol differentially modulates the dynamics of microtubules assembled from unfractionated and purified beta-tubulin isotypes. *Biochemistry* 1997;**36**(12):3554–62.
 99. Lu Q, Luduena RF. Removal of beta III isotype enhances taxol induced microtubule assembly. *Cell Struct Funct* 1993;**18**(3):173–82.
 100. Varna M, Lehmann-Che J, Turpin E, Marangoni E, El-Bouchtaoui M, Jeanne M, et al. P53 dependent cell-cycle arrest triggered by chemotherapy in xenografted breast tumors. *Int J Cancer* 2009;**124**(4):991–7.
 101. Chevillard S, Lebeau J, Pouillart P, de Toma C, Beldjord C, Asselain B, et al. Biological and clinical significance of concurrent p53 gene alterations, MDR1 gene expression, and S-phase fraction analyses in breast cancer patients treated with primary chemotherapy or radiotherapy. *Clin Cancer Res* 1997;**3**(12 Pt 1):2471–8.
 102. Delia D, Mizutani S, Lamorte G, Goi K, Iwata S, Pierotti MA. P53 activity and chemotherapy. *Nat Med* 1996;**2**(7):724–5.
 103. Wattel E, Preudhomme C, Hecquet B, Vanrumbeke M, Quesnel B, Dervite I, et al. P53 mutations are associated with resistance to chemotherapy and short survival in hematologic malignancies. *Blood* 1994;**84**(9):3148–57.
 104. Noguchi S. Predictive factors for response to docetaxel in human breast cancers. *Cancer Sci* 2006;**97**(9):813–20.
 105. Vasey PA, Jones NA, Jenkins S, Dive C, Brown R. Cisplatin, camptothecin, and taxol sensitivities of cells with p53-associated multidrug resistance. *Mol Pharmacol* 1996;**50**(6):1536–40.
 106. Huang Y, Ibrado AM, Reed JC, Bullock G, Ray S, Tang C, et al. Co-expression of several molecular mechanisms of multidrug resistance and their significance for paclitaxel cytotoxicity in human AML HL-60 cells. *Leukemia* 1997;**11**(2):253–7.
 107. Strobel T, Swanson L, Korsmeyer S, Cannistra SA. BAX enhances paclitaxel-induced apoptosis through a p53-independent pathway. *Proc Natl Acad Sci U S A* 1996;**93**(24):14094–9.
 108. Wang Z, Goulet 3rd R, Stanton KJ, Sadaria M, Nakshatri H. Differential effect of anti-apoptotic genes Bcl-xL and c-FLIP on sensitivity of MCF-7 breast cancer cells to paclitaxel and docetaxel. *Anticancer Res* 2005;**25**(3c):2367–79.
 109. Ganta S, Amiji M. Coadministration of Paclitaxel and curcumin in nanoemulsion formulations to overcome multidrug resistance in tumor cells. *Mol Pharm* 2009;**6**(3):928–39.
 110. Mabuchi S, Ohmichi M, Nishio Y, Hayasaka T, Kimura A, Ohta T, et al. Inhibition of inhibitor of nuclear factor-kappaB phosphorylation increases the efficacy of paclitaxel in vitro and in vivo ovarian cancer models. *Clin Cancer Res* 2004;**10**(22):7645–54.
 111. Coloff JL, Macintyre AN, Nichols AG, Liu T, Gallo CA, Plas DR, et al. Akt-dependent glucose metabolism promotes mcl-1 synthesis to maintain cell survival and resistance to bcl-2 inhibition. *Cancer Res* 2011;**71**(15):5204–13.
 112. Bratton MR, Duong BN, Elliott S, Weldon CB, Beckman BS, McLachlan JA, et al. Regulation of ERalpha-mediated transcription of Bcl-2 by PI3K-AKT crosstalk: implications for breast cancer cell survival. *Int J Oncol* 2010;**37**(3):541–50.
 113. Wang X, Yi L, Zhu Y, Zou J, Hong Y, Zheng W. AKT signaling pathway in invasive ductal carcinoma of the breast: correlation with ERa, ERbeta and HER-2 expression. *Tumori* 2011;**97**(2):185–90.
 114. Kneuferrmann C, Lu Y, Liu B, Jin W, Liang K, Wu L, et al. HER2/PI-3K/Akt activation leads to a multidrug resistance in human breast adenocarcinoma cells. *Oncogene* 2003;**22**(21):3205–12.
 115. Sastre-Garau X, Genin P, Rousseau A, Al Ghuzlan A, Nicolas A, Freneau P, et al. Increased cell size and Akt activation in HER-2/neu-overexpressing invasive ductal carcinoma of the breast. *Histopathology* 2004;**45**(2):142–7.
 116. Skoufias DA, Andreassen PR, Lacroix FB, Wilson L, Margolis RL. Mammalian mad2 and bub1/bubR1 recognize distinct spindle-attachment and kinetochore-tension checkpoints. *Proc Natl Acad Sci U S A* 2001;**98**(8):4492–7.
 117. Takahashi T, Yamasaki F, Sudo T, Itamochi H, Adachi S, Tamamori-Adachi M, et al. Cyclin A-associated kinase activity is needed for paclitaxel sensitivity. *Mol Cancer Ther* 2005;**4**(7):1039–46.
 118. Chabalier C, Lamare C, Racca C, Privat M, Valette A, Larminat F. BRCA1 downregulation leads to premature inactivation of spindle checkpoint and confers paclitaxel resistance. *Cell Cycle* 2006;**5**(9):1001–7.
 119. Egawa C, Miyoshi Y, Takamura Y, Taguchi T, Tamaki Y, Noguchi S. Decreased expression of BRCA2 mRNA predicts favorable response to docetaxel in breast cancer. *Int J Cancer* 2001;**95**(4):255–9.
 120. Ali E, Yang JM, Ford JM, Hait WN. Reversal of stathmin-mediated resistance to paclitaxel and vinblastine in human breast carcinoma cells. *Mol Pharmacol* 2007;**71**(5):1233–40.
 121. Chuthapishit S, Layfield R, Kerr ID, Hughes C, Eremin O. Proteomic profiling of MCF-7 breast cancer cells with chemoresistance to different types of anticancer drugs. *Int J Oncol* 2007;**30**(6):1545–51.
 122. Yu D, Jing T, Liu B, Yao J, Tan M, McDonnell TJ, et al. Overexpression of ErbB2 blocks Taxol-induced apoptosis by upregulation of p21Cip1, which inhibits p34Cdc2 kinase. *Mol Cell* 1998;**2**(5):581–91.
 123. Tan M, Jing T, Lan KH, Neal CL, Li P, Lee S, et al. Phosphorylation on tyrosine-15 of p34(Cdc2) by ErbB2 inhibits p34(Cdc2) activation and is involved in resistance to taxol-induced apoptosis. *Mol Cell* 2002;**9**(5):993–1004.
 124. Chang JC, Wooten EC, Tsimelzon A, Hilsenbeck SG, Gutierrez MC, Elledge R, et al. Gene expression profiling for the prediction of therapeutic response to docetaxel in patients with breast cancer. *Lancet* 2003;**362**(9381):362–9.

125. Iwao-Koizumi K, Matoba R, Ueno N, Kim SJ, Ando A, Miyoshi Y, et al. Prediction of docetaxel response in human breast cancer by gene expression profiling. *J Clin Oncol* 2005;**23**(3):422–31.
126. Liu B, Sun D, Xia W, Hung MC, Yu D. Cross-reactivity of C219 anti-p170(mdr-1) antibody with p185(c-erbB2) in breast cancer cells: cautions on evaluating p170(mdr-1). *J Natl Cancer Inst* 1997;**89**(20):1524–9.
127. Mechetner E, Kyshtobayeva A, Zonis S, Kim H, Stroup R, Garcia R, et al. Levels of multidrug resistance (MDR1) P-glycoprotein expression by human breast cancer correlate with in vitro resistance to taxol and doxorubicin. *Clin Cancer Res* 1998;**4**(2):389–98.
128. Trock BJ, Hilakivi-Clarke L, Clarke R. Meta-analysis of soy intake and breast cancer risk. *J Natl Cancer Inst* 2006;**98**(7):459–71.
129. Fujita T, Ito K, Izumi H, Kimura M, Sano M, Nakagomi H, et al. Increased nuclear localization of transcription factor Y-box binding protein 1 accompanied by up-regulation of P-glycoprotein in breast cancer pretreated with paclitaxel. *Clin Cancer Res* 2005;**11**(24 Pt 1):8837–44.
130. Orr GA, Verdier-Pinard P, McDaid H, Horwitz SB. Mechanisms of Taxol resistance related to microtubules. *Oncogene* 2003;**22**(47):7280–95.
131. Kavallaris M. Microtubules and resistance to tubulin-binding agents. *Nat Rev Cancer* 2010;**10**(3):194–204.
132. Berrieman HK, Lind MJ, Cawkwell L. Do beta-tubulin mutations have a role in resistance to chemotherapy? *Lancet Oncol* 2004;**5**(3):158–64.
133. Bernard-Marty C, Treilleux I, Dumontet C, Cardoso F, Fellous A, Gancberg D, et al. Microtubule-associated parameters as predictive markers of docetaxel activity in advanced breast cancer patients: results of a pilot study. *Clin Breast Cancer* 2002;**3**(5):341–5.
134. Hasegawa S, Miyoshi Y, Egawa C, Ishitobi M, Taguchi T, Tamaki Y, et al. Prediction of response to docetaxel by quantitative analysis of class I and III beta-tubulin isotype mRNA expression in human breast cancers. *Clin Cancer Res* 2003;**9**(8):2992–7.
135. Smoter M, Bodnar L, Duchnowska R, Stec R, Grala B, Szczylik C. The role of Tau protein in resistance to paclitaxel. *Cancer Chemother Pharmacol* 2011;**68**(3):553–7.
136. Comen E, Davids M, Kirchhoff T, Hudis C, Offit K, Robson M. Relative contributions of BRCA1 and BRCA2 mutations to “triple-negative” breast cancer in Ashkenazi Women. *Breast Cancer Res Treat* 2011;**129**(1):185–90.
137. Chang HR, Glaspy J, Allison MA, Kass FC, Elashoff R, Chung DU, et al. Differential response of triple-negative breast cancer to a docetaxel and carboplatin-based neoadjuvant treatment. *Cancer* 2010;**116**(18):4227–37.
138. Wang S, Shi Y, Yuan Z, Wang X, Liu D, Peng R, et al. Classical CMF regimen as adjuvant chemotherapy for triple-negative breast cancer may be more effective compared with anthracycline or taxane-based regimens. *Med Oncol* 2011.
139. Wysocki PJ, Korski K, Lamperska K, Zalusk J, Mackiewicz A. Primary resistance to docetaxel-based chemotherapy in metastatic breast cancer patients correlates with a high frequency of BRCA1 mutations. *Med Sci Monit* 2008;**14**(7):SC7–10.
140. Juul N, Szallasi Z, Eklund AC, Li Q, Burrell RA, Gerlinger M, et al. Assessment of an RNA interference screen-derived mitotic and ceramide pathway metagene as a predictor of response to neoadjuvant paclitaxel for primary triple-negative breast cancer: a retrospective analysis of five clinical trials. *Lancet Oncol* 2010;**11**(4):358–65.
141. Yu D. Mechanisms of ErbB2-mediated paclitaxel resistance and trastuzumab-mediated paclitaxel sensitization in ErbB2-overexpressing breast cancers. *Semin Oncol* 2001;**28**(5 Suppl. 16):12–7.
142. Pegram MD, Finn RS, Arzoo K, Beryt M, Pietras RJ, Slamon DJ. The effect of HER-2/neu overexpression on chemotherapeutic drug sensitivity in human breast and ovarian cancer cells. *Oncogene* 1997;**15**(5):537–47.
143. Jarvinen TA, Liu ET. Simultaneous amplification of HER-2 (ERBB2) and topoisomerase IIalpha (TOP2A) genes—molecular basis for combination chemotherapy in cancer. *Curr Cancer Drug Targets* 2006;**6**(7):579–602.
144. Arriola E, Marchio C, Tan DS, Drury SC, Lambros MB, Natrajan R, et al. Genomic analysis of the HER2/TOP2A amplicon in breast cancer and breast cancer cell lines. *Lab Invest* 2008;**88**(5):491–503.
145. Jarvinen TA, Tanner M, Barlund M, Borg A, Isola J. Characterization of topoisomerase II alpha gene amplification and deletion in breast cancer. *Genes Chromosomes Cancer* 1999;**26**(2):142–50.
146. Andre F, Mazouni C, Liedtke C, Kau SW, Frye D, Green M, et al. HER2 expression and efficacy of preoperative paclitaxel/FAC chemotherapy in breast cancer. *Breast Cancer Res Treat* 2008;**108**(2):183–90.
147. Lee JJ, Swain SM. Development of novel chemotherapeutic agents to evade the mechanisms of multidrug resistance (MDR). *Semin Oncol* 2005;**32**:S22–6.
148. Schneider J, Lucas R, Sanchez J, Ruibal A, Tejerina A, Martin M. Modulation of molecular marker expression by induction chemotherapy in locally advanced breast cancer: correlation with the response to therapy and the expression of MDR1 and LRP. *Anticancer Res* 2000;**20**:4373–7.
149. Filipits M, Pohl G, Rudas M, Dietze O, Las X, Grill R, et al. Clinical role of multidrug resistance protein 1 expression in chemotherapy resistance in early-stage breast cancer: the Austrian Breast and Colorectal Cancer Study Group. *J Clin Oncol* 2005;**23**:1161–8.
150. Perez EA. Microtubule inhibitors: differentiating tubulin-inhibiting agents based on mechanisms of action, clinical activity, and resistance. *Mol Cancer Ther* 2008;**8**:2086–95.
151. Rivera E, Fomez H. Chemotherapy resistance in metastatic breast cancer: the evolving role of ixabepilone. *Breast Cancer Res* 2010;**12**(Suppl. 2):S2.
152. Thomas E, Tabern J, Fornier M, Conte P, Fumoleau P, Lluch A, et al. Phase II clinical trial of ixabepilone (BMS-247550), an epothilone B analog, in patients with taxane-resistant metastatic breast cancer. *J Clin Oncol* 2007;**25**:3399–406.
153. Perez EA, Lerzo G, Pivot X, Thomas E, Vahdat L, Bosserman L, et al. Efficacy and safety of ixabepilone (BMS-247550) in a phase II study of patients with advanced breast cancer resistant to an anthracycline, a taxane, and capecitabine. *J Clin Oncol* 2007;**25**:3407–14.
154. Bunnell C, Vahdat L, Schwartzberg L, Gralow J, Klimovsky J, Poulart V, et al. Phase I/II study of ixabepilone plus capecitabine in anthracycline-pretreated/resistant and taxane-resistant metastatic breast cancer. *Clin Breast Cancer* 2008;**8**:234–41.
155. Cleator S, Heller W, Coombes RC. Triple-negative breast cancer: therapeutic options. *Lancet Oncol* 2007;**8**:235–44.
156. Thomas E. Ixabepilone plus capecitabine for metastatic breast cancer progressing after anthracycline and taxane treatment. *J Clin Oncol* 2007;**25**:5210–7.
157. Valero V, Jones SE, Von Hoff DD, Booser DJ, Mennel RG, Ravdin PM, et al. A phase II study of docetaxel in patients with paclitaxel-resistant metastatic breast cancer. *J Clin Oncol* 1998;**16**(10):3362–8.
158. Szakacs G, Paterson JK, Ludwig JA, Booth-Gentle C, Gottesman MM. Targeting multidrug resistance in cancer. *Nat Rev Drug Discov* 2006;**5**(3):219–34.
159. Arai T, Miyoshi Y, Kim SJ, Akazawa K, Maruyama N, Taguchi T, et al. Association of GSTP1 expression with resistance to docetaxel and paclitaxel in human breast cancers. *Eur J Surg Oncol* 2008;**34**(7):734–8.
160. Soussi T, Ishioka C, Claustres M, Beroud C. Locus-specific mutation databases: pitfalls and good practice based on the p53 experience. *Nat Rev Cancer* 2006;**6**(1):83–90.
161. Vincent-Salomon A, Rousseau A, Jouve M, Beuzebec P, Sigal-Zafrani B, Freneaux P, et al. Proliferation markers predictive of the pathological response and disease outcome of patients with breast carcinomas treated by anthracycline-based preoperative chemotherapy. *Eur J Cancer* 2004;**40**(10):1502–8.
162. Daidone MG, Silvestrini R. Prognostic and predictive role of proliferation indices in adjuvant therapy of breast cancer. *J Natl Cancer Inst Monogr* 2001;**30**:27–35.
163. Amadori D, Volpi A, Maltoni R, Nanni O, Amaducci L, Amadori A, et al. Cell proliferation as a predictor of response to chemotherapy in metastatic breast cancer: a prospective study. *Breast Cancer Res Treat* 1997;**43**(1):7–14.
164. Miller K, Wang M, Gralow J, Dickler M, Cobleigh M, Perez EA, et al. Paclitaxel plus bevacizumab versus paclitaxel alone for metastatic breast cancer. *N Engl J Med* 2007;**357**(26):2666–76.
165. Sledge GW, Neuberg D, Bernardo P, Ingle JN, Martino S, Rowinsky EK, et al. Phase III trial of doxorubicin, paclitaxel, and the combination of doxorubicin and paclitaxel as front-line chemotherapy for metastatic breast cancer: an intergroup trial (E1193). *J Clin Oncol* 2003;**21**(4):588–92.
166. Bishop JF, Dewar J, Toner G, Tattersall MH, Oliver I, Ackland S, et al. A randomized study of paclitaxel versus cyclophosphamide/methotrexate/5-fluorouracil/prednisone in previously untreated patients with advanced breast cancer: preliminary results. Taxol Investigational Trials Group, Australia/New Zealand. *Semin Oncol* 1997;**24**(5 Suppl 17): S17–S17-19.
167. Stemmler HJ, Harbeck N, Groll de Rivera I, Vehling-Kaiser U, Rauthe G, Abenhardt W, et al. Prospective multicenter randomized phase III study of weekly versus standard docetaxel plus doxorubicin (D4) for first-line treatment of metastatic breast cancer. *Oncology* 2010;**79**(3–4):204–10.
168. Joensuu H, Sillanpaa L, Alanko T, Sunela K, Huhtanen R, Utraiainen M, et al. Docetaxel versus docetaxel alternating with gemcitabine as treatments of advanced breast cancer: final analysis of a randomised trial. *Ann Oncol* 2010;**21**(5):968–73.
169. Robert N, Leyland-Jones B, Asmar L, et al. Randomized phase III study of trastuzumab, paclitaxel, and carboplatin compared with trastuzumab and paclitaxel in women with HER-2-overexpressing metastatic breast cancer. *J Clin Oncol* 2006;**24**:2786–92.
170. Pegram M, Forbes J, Pienkowski T, et al. BCIRG 007: first overall survival analysis of randomized phase III trial of trastuzumab plus docetaxel with or without carboplatin as first line therapy in HER2 amplified metastatic breast cancer (MBC). *J Clin Oncol* 2007;**25**:968S.
171. Marty M, Cognetti F, Maraninchi D, et al. Superior long-term survival benefits of trastuzumab plus docetaxel compared to docetaxel alone in patients with HER2-positive metastatic breast cancer: patients surviving more than 4 years in the M77001 study. *Breast Cancer Res Treat* 2006;**100**(Suppl. 1):S103.
172. Kim SJ, Miyoshi Y, Taguchi T, Tamaki Y, Nakamura H, Yodoi J, et al. High thioredoxin expression is associated with resistance to docetaxel in primary breast cancer. *Clin Cancer Res* 2005;**11**(23):8425–30.
173. Martin M, Romero A, Cheang MC, Lopez Garcia-Asenjo JA, Garcia-Saenz JA, Oliva B, et al. Genomic predictors of response to doxorubicin versus docetaxel in primary breast cancer. *Breast Cancer Res Treat* 2011;**128**(1):127–36.
174. Sankaran S, Starita LM, Groen AC, Ko MJ, Parvin JD. Centrosomal microtubule nucleation activity is inhibited by BRCA1-dependent ubiquitination. *Mol Cell Biol* 2005;**25**(19):8656–68.
175. Estevez LG, Cuevas JM, Anton A, Florian J, Lopez-Vega JM, Velasco A, et al. Weekly docetaxel as neoadjuvant chemotherapy for stage II and III breast cancer: efficacy and correlation with biological markers in a phase II, multicenter study. *Clin Cancer Res* 2003;**9**(2):686–92.

176. Miyoshi Y, Kim SJ, Akazawa K, Kamigaki S, Ueda S, Yanagisawa T, et al. Down-regulation of intratumoral aromatase messenger RNA levels by docetaxel in human breast cancers. *Clin Cancer Res* 2004;**10**(24):8163–9.
177. Modi S, DiGiovanna MP, Lu Z, Moskowitz C, Panageas KS, Van Poznak C, et al. Phosphorylated/activated HER2 as a marker of clinical resistance to single agent taxane chemotherapy for metastatic breast cancer. *Cancer Invest* 2005;**23**(6):483–7.
178. Sjostrom J, Collan J, von Boguslawski K, Franssila K, Bengtsson NO, Mjaaland I, et al. C-erbB-2 expression does not predict response to docetaxel or sequential methotrexate and 5-fluorouracil in advanced breast cancer. *Eur J Cancer* 2002;**38**(4):535–42.
179. Gonzalez-Angulo AM, Krishnamurthy S, Yamamura Y, Broglio KR, Pusztai L, Buzdar AU, et al. Lack of association between amplification of her-2 and response to preoperative taxanes in patients with breast carcinoma. *Cancer* 2004;**101**(2):258–63.
180. Camerini A, Donati S, Viacava P, Siclari O, Puccetti C, Tartarelli G, et al. Evaluation of HER2 and p53 expression in predicting response to docetaxel-based first-line chemotherapy in advanced breast cancer. *J Exp Clin Cancer Res* 2011;**30**:38.
181. Di Leo A, Chan S, Paesmans M, Friedrichs K, Pinter T, Cocquyt V, et al. HER-2/neu as a predictive marker in a population of advanced breast cancer patients randomly treated either with single-agent doxorubicin or single-agent docetaxel. *Breast Cancer Res Treat* 2004;**86**(3):197–206.
182. Luftner D, Henschke P, Flath B, Akrivakis C, Schnabel S, Prinz B, et al. Serum HER-2/neu as a prediction and monitoring parameter in a phase II study with weekly paclitaxel in metastatic breast cancer. *Anticancer Res* 2004;**24**(2B):895–906.
183. Ooe A, Kato K, Noguchi S. Possible involvement of CCT5, RGS3, and YKT6 genes up-regulated in p53-mutated tumors in resistance to docetaxel in human breast cancers. *Breast Cancer Res Treat* 2007;**101**(3):305–15.
184. Chrisanthar R, Knappskog S, Lokkevick E, Anker G, Ostenstad B, Lundgren S, et al. Predictive and prognostic impact of TP53 mutations and MDM2 promoter genotype in primary breast cancer patients treated with epirubicin or paclitaxel. *PLoS One* 2011;**6**(4):e19249.
185. Maeno K, Ito K, Hama Y, Shingu K, Kimura M, Sano M, et al. Mutation of the class I beta-tubulin gene does not predict response to paclitaxel for breast cancer. *Cancer Lett* 2003;**198**(1):89–97.